arXiv:1606.03837v1 [math.NT] 13 Jun 2016

Identities on the k-ary Lyndon words related to a family of zeta functions

Irem Kucukoglu[1,a] and Yilmaz Simsek[1,b]
[a]ikucukoglu@akdeniz.edu.tr
[b]ysimsek@akdeniz.edu.tr
[1]Department of Mathematics, Faculty of Science University of Akdeniz
TR-07058 Antalya, Turkey

Abstract

The main aim of this paper is to investigate and introduce relations between the numbers of k-ary Lyndon words and unified zeta-type functions which was defined by Ozden et al.[15, p. 2785, Definition 3]. Finally, we give some identities on generating functions for the numbers of k-ary Lyndon words and some special numbers and polynomials such as the Apostol-Bernoulli numbers and polynomials, Frobenius-Euler numbers, Euler numbers and Bernoulli numbers.

2010 Mathematics Subject Classification: 03D40, 05A05, 05A15, 11A25, 11B68, 11B83, 11F22, 11S40, 11M99, 68R15, 94B40.

Keywords: Lyndon words, Generating functions, Special numbers, Special polynomials, Arithmetical functions, Hurwitz-Lerch zeta functions, Apostol-Bernoulli numbers and polynomials, Frobenius-Euler numbers, Euler numbers, Bernoulli numbers.

1 Introduction

Throughout this paper, we consider the number of k-ary Lyndon words of length n, $L_k(n)$ as follows [4]:

$$L_k(n) = \frac{1}{n}\sum_{d|n}\mu\left(\frac{n}{d}\right)k^d, \tag{1}$$

where the arithmetic function μ is the Möbius function defined as follows [2]:

$$\mu(1) = 1;$$

Let p_1, p_1,..., p_k be k distinct primes. If $n > 1$, write $n = p_1^{a_1}p_2^{a_2}\dots p_k^{a_k}$. Then

$$\begin{aligned}\mu(n) &= (-1)^k \text{ if } a_1 = a_2 = \dots = a_k = 1,\\ &\quad \text{that is, if } n \text{ is the product of } k \text{ distinct primes,}\\ \mu(n) &= 0 \text{ if } n \text{ is the product of non distinct primes.}\end{aligned}$$

In [12], Lyndon words are studied as standard lexicographic sequences. According to [4, p. 36], *a k-ary necklace is an equivalence class of k-ary strings under rotation. As a representative of such an equivalence class which is taken the smallest in the lexicographical order. A period n necklace representative with n digits is called a Lyndon word.*

In addition to counting k-ary Lyndon words of length n, Equation (1) is well known as Witt's formula which is used to count the number of monic irreducible polynomials of degree n over Galois field (*cf.* [3]) and As we mentioned previously in [10], it is know that there are really interesting connections between this formula and dimension formula for the homogeneous subspaces of the free Lie algebra (*cf.* [3], [8], [18]) and the rank of the free abelian quotient (*cf.* [29], [12]). Furthermore, it is also called necklace polynomial (*cf.* [13]). For further information about $L_k(n)$ and table including numerical values of $L_k(n)$, the reader can consult [10], its references and also the references cited in each of these earlier works.

In [10], the authors gave the following explicit formula for the numbers of $L_k(p)$ for p is a prime number, $m \in \mathbb{N}$:

$$L_k(p^m) = \frac{k^{p^{m-1}}\left(k^{p^{m-1}(p-1)} - 1\right)}{p^m}. \tag{2}$$

In [10], the authors defined ordinary generating functions for the numbers of k-ary Lyndon words of prime length p, $L_k(p)$ for prime p, as follows:

Let p is a prime number and $m = 1$ in the special case of Equation (2)

$$\mathfrak{f}_L(t,p) = \sum_{k=2}^{\infty} L_k(p)\, t^k = \frac{1}{p}\sum_{k=2}^{\infty}\left(k^p - k\right) t^k, \tag{3}$$

(*cf.* [10]). However, we modify (3) as follows:

$$f_{L_y}(t,p) = \sum_{k=0}^{\infty} L_k(p)\, t^k.$$

2 Relation between $f_{L_y}(t,p)$ and a family of zeta functions

In this section, our aim is to give some identities on the generating functions for the numbers of k-ary Lyndon words of length prime related a family of zeta-type function, the Apostol-Bernoulli numbers and polynomials, Frobenius-Euler numbers and Euler numbers.

The Apostol-Bernoulli numbers, $\mathcal{B}_k(z)$ are defined by means of the following generating functions:

$$\frac{t}{ze^t - 1} = \sum_{k=0}^{\infty} \frac{\mathcal{B}_k(z)}{k!} t^k,$$

(*cf. [1]*). The Apostol Benoulli polynomials, $\mathcal{B}_k(x,z)$ are also defined by means of the following generating functions:

$$\frac{te^{tx}}{ze^t - 1} = \sum_{k=0}^{\infty} \frac{\mathcal{B}_k(x,z)}{k!} t^k,$$

where $|t| < 2\pi$ when $z = 1$ and $|t| < |\log z|$ when $z \neq 1$ and $z \in \mathbb{C}$. From this generating function, one can easily see that

$$\mathcal{B}_m(x,z) = \sum_{j=0}^{m} \binom{m}{j} x_j^{m-j} \mathcal{B}_j(z),$$

and

$$\mathcal{B}_m(0,z) = \mathcal{B}_m(z).$$

By using the above generating function, several of Apostol Bernoulli numbers and polynomials are given as follows, respectively (*cf. [1]*):

$$\begin{aligned}
\mathcal{B}_0(z) &= 0, \\
\mathcal{B}_1(z) &= \frac{1}{z-1}, \\
\mathcal{B}_2(z) &= \frac{-2z}{(z-1)^2}, \\
\mathcal{B}_3(z) &= \frac{3z(z+1)}{(z-1)^3}, \\
\mathcal{B}_4(z) &= \frac{-4z\left(z^2+4z+1\right)}{(z-1)^4}, \\
\mathcal{B}_5(z) &= \frac{5z\left(z^3+11z^2+11z+1\right)}{(z-1)^5}, \\
\mathcal{B}_6(z) &= \frac{-6z\left(z^4+26z^3+66z^2+26z+1\right)}{(z-1)^6}
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{B}_0(z,x) &= 0, \\
\mathcal{B}_1(z,x) &= \frac{1}{z-1}, \\
\mathcal{B}_2(z,x) &= \frac{1}{z-1}x - \frac{2z}{(z-1)^2}, \\
\mathcal{B}_3(z,x) &= \frac{9z(z-1)}{(z-1)^3}x^2 - \frac{6z}{(z-1)^2}x + \frac{3z(z-1)}{(z-1)^3},
\end{aligned}$$

(*cf. [1], [6], [15], [16], [22], [26], [27], [28]*).

In the above numerical computation of the Apostol-Bernoulli numbers, we observe that all of this numbers are rational functions of parameter z. $z = 1$ is a pole of these functions.

The following generating function of the unification of the Bernoulli, Euler and Genocchi polynomials, $\mathcal{Y}_{n,\beta}(x;k,a,b)$, which was recently defined by Ozden [14] for $k \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} (\mathbb{N} := \{1,2,3,\ldots\})$; $a,b \in \mathbb{R}^+$; $\beta \in \mathbb{C}$:

$$\frac{2^{1-k}t^k e^{tx}}{\beta^b e^t - a^b} = \sum_{n=0}^{\infty} \mathcal{Y}_{n,\beta}(x;k,a,b)\frac{t^n}{n!} \tag{4}$$

where $\left|t + b\log\left(\frac{\beta}{a}\right)\right| < 2\pi$; $x \in \mathbb{R}$ and note that

$$\mathcal{Y}_{n,\beta}(k,a,b) = \mathcal{Y}_{n,\beta}(0;k,a,b) = \mathcal{Y}_{n,\beta}(1;k,a,b).$$

The following equation of the unified zeta-type functions $\zeta_\beta(s,x;k,a,b)$, which was recently defined by Ozden et al.[15, p. 2785, Definition 3]:

$$\zeta_\beta(s,x;k,a,b) = \left(\frac{-1}{2}\right)^{k-1} \sum_{n=0}^{\infty} \frac{\beta^{bn}}{a^{b(n+1)}(n+x)^s} \tag{5}$$

where $\beta, s \in \mathbb{C}$ with $Re(s) < 1$ and $|\beta| < 1$ and observe that if $x = 1$, then

$$\zeta_\beta(s:k,a,b) = \zeta_\beta(s,1;k,a,b) = \left(\frac{-1}{2}\right)^{k-1} \sum_{n=1}^{\infty} \frac{\beta^{b(n-1)}}{a^{b(n+1)}n^s}.$$

Ozden et al.[15, p. 2789, Theorem 7] also proved the following relation for $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$:

$$\zeta_\beta(1-n:x;a,b) = (-1)^k \frac{(n-1)!}{(n+k-1)!}\mathcal{Y}_{n+k-1,\beta}(x;k,a,b). \tag{6}$$

Remark 1 *Setting $s \to -m$, $\beta \to t$, $x = 0$, $k = a = b = 1$ in (5), we have*

$$\zeta_\beta(-m,0;1,1,1) = \sum_{n=0}^{\infty} t^n n^s \tag{7}$$

and also setting $1 - n \to -m$, $\beta \to t$, $x = 0$, $k = a = b = 1$ in (6), we get

$$\zeta_t(-m:0;1,1) = -\frac{\mathcal{Y}_{1+m,t}(0;1,1,1)}{1+m} \tag{8}$$

It is well-know that $\mathcal{Y}_{n,\beta}(x;1,1,1)$ reduce to the Apostol-Bernoulli polynomials, $\mathcal{B}_n(x,\beta)$ and Apostol-Bernoulli numbers $\mathcal{B}_n(\beta)$, respectively. Thus

$$\mathcal{Y}_{n,\beta}(0;1,1,1) = \mathcal{B}_n(0,\beta) = \mathcal{B}_n(\beta). \tag{9}$$

Hurwitz-Lerch zeta function $\Phi(z,s,a)$ is defined by (*cf.* [28, p. 121 et seq.], [28, p. 194 et seq.]):

$$\Phi(z,s,a) = \sum_{n=0}^{\infty} \frac{z^n}{(n+a)^s},$$

where $a \in \mathbb{C}\backslash\mathbb{Z}_0^-$; s $\in \mathbb{C}$ when $|z| < 1$; $Re(s) > 1$ when $|z| = 1$. One can also easily see that a relation between $\zeta_\beta(s,x;k,a,b)$ and $\Phi(z,s,a)$ is given by

$$\zeta_\beta(s,x;k,a,b) = \frac{\left(\frac{-1}{2}\right)^{k-1}}{a^b}\Phi\left(\frac{\beta^b}{a^b},s,x\right)$$

(*cf.* [15, Eq-(4.14)]). Hurwitz–Lerch zeta function is related to not only Riemann zeta function and the Hurwitz zeta function (see, for details, [28, Chapter 2], see also [27], [22]):

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \Phi(1, s, 1),$$

and

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s} = \Phi(1, s, a).$$

Remark 2 *Let $n \geq 1$. Then in [9], [23] and [24], one can see that*

$$\mathcal{B}_n(z) = \frac{n}{z^{-1}} \mathcal{H}_{n-1}\left(\frac{1}{z}\right), \tag{10}$$

where $\mathcal{H}_n(z)$ denotes the Frobenius-Euler numbers which are defined by means of the following generating function:

$$\frac{1-z}{e^t - z} = \sum_{n=0}^{\infty} \mathcal{H}_n(z) \frac{t^n}{n!},$$

for $z = -1$, we have $\mathcal{H}_n(-1) = E_n$ which is defined by

$$\frac{2}{e^t + 1} = \sum_{n=0}^{\infty} E_n(z) \frac{t^n}{n!},$$

(cf. [7]-[28]; and the references cited therein).

Now, by combining Equation (7) with Equation (8) and Equation (9), we obtain the following explicit formula of generating functions for the numbers of k-ary Lyndon words in terms of the Apostol-Bernoulli numbers:

Theorem 3 *Let p is a prime number. Then*

$$f_{L_y}(t, p) = \frac{\mathcal{B}_2(t)}{2p} - \frac{\mathcal{B}_{p+1}(t)}{p(p+1)}$$

where $\mathcal{B}_{p+1}(z)$ denotes the Apostol-Bernoulli numbers.

Remark 4 *If we substitute $p = 2$ into Theorem 3, we arrive at*

$$f_{L_y}(t, 3) = \frac{t^2}{(1-t)^3}$$

which was given in [10, p. 3].

Remark 5 *Observe that degree of polynomial in the numerator of $f_{L_y}(t,2)$ is lower than its the denominator. Also, when $p=3$ into Theorem 3, we also arrive at*

$$f_{L_y}(t,3)=\frac{2t^2}{(t-1)^4}.$$

By combining Theorem 3 with Equation (10), we give the following Remark:

Theorem 6 *Let p is a prime number. Then*

$$f_{L_y}(t,p)=\frac{t\left(\mathcal{H}_1\left(\frac{1}{t}\right)-\mathcal{H}_p\left(\frac{1}{t}\right)\right)}{p} \tag{11}$$

where $\mathcal{H}_p\left(\frac{1}{t}\right)$ denotes the Frobenius-Euler number.

Since $\mathcal{H}_n(-1)=E_n$, then Equeation (11) is reduced to the following Corollary:

Corollary 7

$$f_{L_y}(-1,p)=\frac{E_p-E_1}{p}$$

where E_n denotes Euler numbers.

3 Further identities related to Bernoulli numbers

In this section, we give some identities on the numbers of k-ary Lyndon words related to the Apostol-Bernoulli numbers. Now, we recall definition the Bernoulli polynomials $B_n(x)$ which are defined by means of the following generating function:

$$\frac{te^{tx}}{e^t-1}=\sum_{n=0}^{\infty}B_n(x)\frac{t^n}{n!},$$

where $|t|<2\pi$ and also

$$B_n=B_n(0)$$

which denotes the Bernoulli numbers (*cf.* [7]-[28]; and the references cited therein).

The sum of the powers of integers is related to the Bernoulli numbers and polynomials:

$$\sum_{k=0}^{m}k^n=\frac{B_{n+1}(m+1)-B_{n+1}}{n+1} \tag{12}$$

(*cf.* [5], [28], [25]; see also the references cited in each of these earlier works).

After applying mobius inversion formula to Equation (3), we have

$$k^n=\sum_{d|n}dL_k(d). \tag{13}$$

Hence, summing Equation (13) over all $0 \leq k \leq m$ and combining with Equation (12), we obtain the following Theorem:

Theorem 8 *Let $n \geq 1$. Then*

$$\sum_{d|n} d \sum_{k=0}^{m} L_k(d) = \frac{B_{n+1}(m+1) - B_{n+1}}{n+1},$$

where $B_{n+1}(m)$ denotes Bernoulli polynomials.

Acknowledgments. The present investigation was supported by *Scientific Research Project Administration of Akdeniz University.*

References

[1] T. M. Apostol, On the Lerch zeta function, *Pacific Journal of Mathematics* **1 (**1951), pp. 161-167.

[2] T. M. Apostol, *Introduction to Analytic Number Theory,* Narosa Publishing, Springer Verlag, New Delhi, Chennai, Mumbai, 1998.

[3] H.L. Buchanan, A. Knopfmacher, M.E. Mays, On the cyclotomic identity and related product expansions, *Australas. J. Combin.* **8** (1993), pp. 233-245.

[4] T. W. Cusick and P. Stanica, *Cryptographic Boolean Functions and Applications*, London: Academic Press, Elsevier, 2009.

[5] G. B. Djordjevic and G. V. Milovanovic, Special classes of polynomials, University of Nis, Faculty of Technology Leskovac, 2014.

[6] S. Hu, M.-S. Kim, Two closed forms for the Apostol-Bernoulli polynomials, arXiv:1509.04190.

[7] **L. C. Jang and H. K. Pak, Non-archimedean integration associated with q-Bernoulli numbers, Proc. Jangjeon Math. Soc. 5(2) (2002), pp. 125-129.**

[8] S.-J. Kang, M.-H. Kim. 1996. Free Lie algebras, generalized Witt formula, and the denominator identity. *J. Algebra* **183** (1996), no. 2, pp. 560-594.

[9] **T. Kim, S.-H. Rim, Y. Simsek, and D Kim, On the analogs of Bernoulli and Euler numbers, related identities and zeta and l-functions, J. Korean Math. Soc. 45(2) (2008), pp. 435-453.**

[10] I. Kucukoglu, Y. Simsek, On k-ary Lyndon Words And Their Generating Functions, to appear in *AIP Conf. Proc. of ICNAAM*(2016).

[11] **Q.-M. Luo, H. M. Srivastava, Some generalizations of the Apostol-Bernoulli and Apostol-Euler polynomials, J. Math. Anal. Appl. 308 (2005), pp. 290-302.**

[12] R. Lyndon, On Burnside problem I, Trans. *American Math. Soc.* **77** (1954), pp. 202-215.

[13] N. Metropolis, G.-C. Rota, Witt Vectors and the Algebra of Necklaces, *Adv. in Math.* **50** (1983), pp. 95-125.

[14] H. Ozden, Unification of generating function of the Bernoulli, Euler and Genocchi numbers and polynomials, *Amer. Inst. Phys. Conf. Proc.* **1281** (2010), 1125-1128.

[15] H. Ozden, Y. Simsek, H.M. Srivastava, A unified presentation of the generating functions of the generalized Bernoulli, Euler and Genocchi polynomials, *Comput. Math. Appl.* **60** (2010), pp. 2779-2787.

[16] H. Ozden, Y. Simsek, Unified presentation of p-adic L-functions associated with unification of the special numbers, *Acta Math. Hungar.* **144** (**2**) (2014), pp. 515-529.

[17] **H. Ozden, Y. Simsek, Modification And Unification Of The Apostol-Type Numbers And Polynomials And Their Applications, *Appl. Math. Comput.* 235 (2014), pp. 338-351.**

[18] V.M. Petrogradsky, Witt's formula for restricted Lie algebras, *Adv. Appl. Math.* **30** (2003), pp. 219-227.

[19] **Y. Simsek, T. Kim, D.W. Park, Y.S. Ro, L.C. Jang, S. Rim, An Explicit Formula For The Multiple Frobenius-Euler Numbers And Polynomials, *JP J. Algebra Number Theory Appl.* (2004), no.3, pp. 519-529.**

[20] **Y. Simsek, q-Analogue of the twisted l-Series and q-Twisted Euler Numbers, *J. Number Theory* 100(2) (2005), pp. 267-278.**

[21] **Y. Simsek, O. Yurekli, V. Kurt, On interpolation functions of the twisted generalized Frobenius–Euler numbers, *Adv. Stud. Contemp. Math.* 15 (2007), no. 2, pp. 187-194.**

[22] Y. Simsek and H. M. Srivastava, A family of p-adic twisted interpolation functions associated with the modified Bernoulli numbers, *Appl. Math. Comput.* **216** (2010), pp. 2976-2987.

[23] **Y. Simsek, Generating functions for generalized Stirling type numbers, Array type polynomials, Eulerian type polynomials and their applications, *Fixed Point Theory A.* 2013, 2013:87**.

[24] **Y. Simsek, Apostol type Dahee numbers and Polynomials, to appear in Adv. Stud. Contemp. Math. 26(3) (2016).**

[25] H. M. Srivastava and H. L. Manocha, *A Treatise on Generating Functions*, Ellis Horwood Limited Publisher, Chichester, 1984.

[26] H.M. Srivastava, J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers, Dordrecht, Boston and London, 2001.

[27] H. M. Srivastava, H. Ozden, I. N. Cangul and Y. Simsek, A unified presentation of certain meromorphic functions related to the families of the partial zeta type functions and the L-functions, *Appl. Math. Comput.* **219** (2012), pp. 3903-3913.

[28] H.M. Srivastava, J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

[29] E. Witt, Treue Darstellung Liescher Ringe, *J. Reine Angew. Math.* **177** (1937), pp.152-160.